



Design and deployment of semiochemical traps for capturing *Anthonomus rubi* Herbst (Coleoptera: Curculionidae) and *Lygus rugulipennis* Poppius (Heteroptera: Miridae) in soft fruit crops

Fountain, Michelle T.; Baroffio, Catherine; Borg-Karlson, Anna-Karin; Brain, Phil; Cross, Jerry V.; Farman, Dudley I.; Hall, David R.; Ralle, Baiba; Rendina, Paulo; Richoz, Pauline; Sigsgaard, Lene; Storberget, Sverre; Trandem, Nina; Wibe, Atle

Published in:
Crop Protection

DOI:
[10.1016/j.cropro.2017.05.001](https://doi.org/10.1016/j.cropro.2017.05.001)

Publication date:
2017

Document version
Peer reviewed version

Document license:
[CC BY-NC-ND](#)

Citation for published version (APA):
Fountain, M. T., Baroffio, C., Borg-Karlson, A-K., Brain, P., Cross, J. V., Farman, D. I., ... Wibe, A. (2017). Design and deployment of semiochemical traps for capturing *Anthonomus rubi* Herbst (Coleoptera: Curculionidae) and *Lygus rugulipennis* Poppius (Heteroptera: Miridae) in soft fruit crops. *Crop Protection*, 99, 1-9. <https://doi.org/10.1016/j.cropro.2017.05.001>

Design and deployment of semiochemical traps for capturing *Anthonomus rubi* Herbst (Coleoptera: Curculionidae) and *Lygus rugulipennis* Poppius (Heteroptera: Miridae) in soft fruit crops

Michelle T. Fountain¹, Catherine Baroffio², Anna-Karin Borg-Karlson⁶, Phil Brain¹, Jerry V. Cross¹, Dudley I. Farman⁷, David R. Hall⁷, Baiba Ralle⁴, Paulo Rendina², Pauline Richoz², Lene Sigsgaard⁵, Sverre Storberget³, Nina Trandem^{2,3} and Atle Wibe⁸

¹NIAB EMR, New Road, East Malling, Kent, ME19 6BJ, UK. Michelle.fountain@emr.ac.uk

²Norwegian Institute of Bioeconomy Research (NIBIO), P.O. Box 115, NO-1431 Ås, Norway

³Department of Ecology and Natural Resource Management, Norwegian University of Life Sciences, P.O. Box 5003, NO-1432 Ås, Norway

²Agroscope IPS, Research Center Conthey, Route des Eterpys 18, 1964 Conthey, Switzerland.

⁴Latvian Plant Protection Research Centre, Struktoru iela 14a Riga LV-1039, Latvia.

⁵University of Copenhagen, Department of Plant and Environmental Sciences, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark.

⁶KTH Royal Institute of Technology, Department of Chemistry, 10044 Stockholm, Sweden.

⁷Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent, ME4 4TB, UK.

⁸NORSØK Norwegian Centre for Organic Agriculture, Gunnars vei 6, NO-6630 Tingvoll, Norway

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

Abstract

Strawberry blossom weevil (SBW), *Anthonomus rubi* Herbst (Coleoptera: Curculionidae) and European tarnished plant bug (ETB), *Lygus rugulipennis* Poppius (Heteroptera: Miridae), cause significant damage to strawberry and raspberry crops. Using the SBW aggregation pheromone and ETB sex pheromone we optimized and tested a single trap for both species. A series of field experiments in crops and semi-natural habitats in five European countries tested capture of the target pests and the ability to avoid captures of beneficial arthropods. A Unitrap containing a trapping agent of water and detergent and with a cross vane was more efficient at capturing both species compared to traps which incorporated glue as a trapping agent. Adding a green cross vane deterred attraction of non-pest species such as bees, but did not compromise catches of the target pests. The trap caught higher numbers of ETB and SBW if deployed at ground level and although a cross vane was not important for catches of ETB it was needed for significant captures of SBW. The potential for mass trapping SBW and ETB simultaneously in soft fruit crops is discussed including potential improvements to make this more effective and economic to deploy.

Key words: Apoidea, bycatch, monitoring, mass trapping, pheromone, plant volatiles

Introduction

Across Europe, strawberry blossom weevil (SBW), *Anthonomus rubi* Herbst (Coleoptera: Curculionidae) and European tarnished plant bug (ETB), *Lygus rugulipennis* Poppius (Heteroptera: Miridae) are serious pests in strawberry and some cane fruits causing economic loss for farmers. SBW females lay eggs in flower buds and then partially sever the peduncles. Damaged buds do not develop further resulting in a loss of yield (Aasen & Trandem, 2006; Jay *et al.*, 2008). ETB pierces and feeds on flowers and developing fruitlets, causing fruit distortion and considerably decreasing fruit quality for market, up to 80% distorted fruits (Cross *et al.*, 2011; Fitzgerald & Jay, 2011).

Foliar applications of insecticides are the main method of controlling these pests. The loss of active compounds through the pesticides approval process, the evolution of pesticide resistance in many pest populations (e.g. in SWB, Aasen & Trandem, 2006), the need for selective control measures to prevent disruption of integrated pest management (IPM) practices (Hillocks, 2012; 2013) and high losses in organic production all require better timed and targeted control applications and alternative control methods for key pest species. In addition, the incidence of pesticide residues in fresh produce (European Food Safety Authority, 2015) and harm to beneficial insects (e.g. Croft & Brown, 1975; Cressey, 2015) are all justifications for alternative approaches to pesticide use (Hillocks, 2012, 2013).

In the EU, users of pesticides are required by law to monitor pests when possible, and only apply pesticides when pests are present in damaging numbers and other measures have failed, taking the resistance risk into account (Sustainable Use Directive 2009/128/EC). The use of pheromone traps for monitoring insect pests is widespread in Europe and other main fruit growing regions of the world (Walton *et al.*, 2004; Teixeira *et al.*, 2009; Haghani *et al.*, 2016). Trap design, placement and attractants may all have an important role in pheromone trap effectiveness, depending on pest behaviour and finding the best combination of these factors will improve trap efficacy (Blackmer *et al.*, 2008; Switzer *et al.*, 2009; Singh *et al.*, 2013; Renkema *et al.*, 2014).

Effective monitoring traps also have the potential to control pests through mass trapping (Faccoli and Stergulc, 2008; Witzgall *et al.*, 2010; Abbes *et al.*, 2012; Mwatawala *et al.*, 2015) aiming to reduce pest numbers, sufficiently, to reduce fruit damage. Mass trapping has been used in the long term management of many pests and has the potential to be exploited for commercial strawberry production by suppressing or even eradicating low-density, isolated pest populations (El-Sayed *et al.*, 2006). The combination of mass trapping and releases of the predator *Nesidiocoris tenuis* (Reuter) resulted in a 50% reduction in tomato fruit infestation by the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), compared to conventional treatments (Abbes *et al.*, 2012). Mass trapping often reduces populations of pests in crops (e.g. Mafra-Neto & Habib, 2003), but there are fewer studies demonstrating successful damage reduction. Examples of successful use of mass trapping against Coleoptera include the spruce bark beetle, *Ips typographus* (L.) (Faccoli & Stergulc, 2008), and the palm weevils *Rhynchophorus palmarum* (L.) (Oehschlager *et al.*, 2002) and *R. ferrugineus* (Olivier) (Dembilio & Jaques, 2015).

The male-produced aggregation pheromone of SBW was identified as a blend of Grandlure I, Grandlure II and lavandulol by Innocenzi *et al.* (2001), and further work was carried out to make the blend more cost-effective by Innocenzi *et al.* (2001) and Cross *et al.* (2006b). In addition, the effect of host plant volatiles on SBW was investigated. Bichão *et al.* (2005a,b) showed that some neurons on the antenna of *A. rubi* are narrowly tuned to a few structurally related sesquiterpenes, aromatics or monoterpenes. Adding these plant volatiles to the aggregation pheromone has the potential to increase the attractiveness to SBW (Cross *et al.*, 2006b; Wibe *et al.*, 2011; 2014a). Currently a blend of SBW aggregation pheromone and one plant volatile, 1,4-dimethoxybenzene, is widely used for SBW monitoring (Wibe *et al.*, 2011; 2014a).

Three compounds have been identified as components of the ETB female sex pheromone (Innocenzi *et al.*, 2004; Frati *et al.*, 2009) and a blend of these was further optimised and tested in field trials (Innocenzi *et al.*, 2004, 2005; Fountain *et al.*, 2008, 2011; Cross *et al.*, 2011) to develop an effective lure and trap for monitoring males (Fountain *et al.*, 2014). In addition, some plant volatiles

such as phenylacetaldehyde have been identified as attractants for female ETB (Fрати *et al.*, 2009; Fountain *et al.*, 2010; Koczor *et al.*, 2012).

For both target species, initial testing assessed different trap types and colours, most frequently using traps which incorporated sticky glue as the trapping agent (Innocenzi *et al.*, 2001; Cross *et al.*, 2006a, 2006b; Jay *et al.*, 2008). These traps were not optimal for SBW as weevils were often found around the traps, but not in or on them (Cross *et al.*, 2006a). Initial experiments for attracting ETB employed various sticky trap designs and colours but this was before the pheromone was widely available (Holopainen *et al.*, 2001; Blackmer *et al.*, 2008).

Changes in trap design leading to improved pest capture will make a monitoring trap more sensitive and mass trapping more effective. Traps must be competitive with the surrounding crop, ensure the pest is captured and not kill or disrupt significant numbers of natural enemies and other beneficial insects, e.g. pollinators. In addition, it should not become saturated with bycatch and it should be easy to use and maintain, and be cost effective.

To help reduce pesticide inputs, further development of the traps was necessary to a) improve target pest capture, b) combine traps for two common species in strawberry and c) develop a trap which was easy to maintain and economically viable for future mass trapping. Studies were carried out in Denmark, Latvia, Norway, Switzerland and the UK comparing the effect of various trap designs on captures of the target pests including non-target, beneficial, species.

Materials and methods

Traps

Two basic designs of trap were evaluated; delta traps (20 cm x 20 cm) with white sticky inserts and green Unitraps consisting of a bucket with a funnelled entrance and green or white cross vanes between the bucket and the roof (bucket 16 cm dia, 12.5 cm high with 3 cm dia opening, cross vanes 10 cm high, cover 16.5 cm dia). The latter trap, from hereon in, will be referred to as Unitraps. Water (250 ml) and a drop of detergent was added to the Unitraps as killing agent. Traps were purchased

from Agrisense (Treforest, Pontypridd, UK), International Pheromone Systems Ltd. (The Wirral, Merseyside, UK) or Agralan (Swindon, UK).

Lures

For trapping ETB with live females, individual mature, virgin, female ETB from a laboratory culture were contained in a cage (hair roller 6 cm x 3 cm with gauze around the outside and a lid at either end, holding the gauze in place). The cage contained a piece of damp paper and a section of bean as food and was anchored into the top of the trap under the roof. Female ETB were replaced weekly.

Lures for SBW were polyethylene sachets containing 100 µl of 1:4:1 blend of Grandlure I: Grandlure II: lavandulol plus 200 mg 1,4-dimethoxybenzene (Wibe *et al.*, 2014) (International Pheromone Systems Ltd.). Lures for ETB were pipette tips containing 10 µg hexyl butyrate, 0.3 µg (*E*)-2-hexenyl butyrate and 2 µg (*E*)-4-oxo-2-hexenal in 100 µl sunflower oil (Fountain *et al.*, 2014), prepared at the Natural Resources Institute. Lures were hung from the roof of delta traps or the cover of Unitraps.

Comparison of delta traps and Unitraps for trapping ETB

Two experiments were carried out in a weed field (*Chenopodium* and *Matricaria*) at NIAB EMR in the UK (Lat: 51.285494 north, Long: 0.461177 east) using virgin female ETB as bait (Table 1). In Experiment A (27 June – 11 July 2008), delta traps and Unitraps were compared with different materials for retaining the insects. The delta traps had either the standard wet glue inserts, dry glue inserts (Agrisense), wet glue inserts with additional sticker or wet glue inserts sprayed with cypermethrin (0.0014 ml sticky base⁻¹, equivalent to 0.35 L ha⁻¹). The Unitraps had white cross vanes or cross vanes constructed from white insect trapping cards impregnated with lambda-cyhalothrin. A clear delta trap was also tested, made of clear vinyl sheets held together at the top with a paper binder and with a white, wet, glue insert (Table 1).

In Experiment B (27 August – 1 September 2008), different coloured traps were compared. These were green delta traps, clear delta traps, and green Unitraps with white, green or yellow cross vanes. A sticky stake trap was also tested consisting of a wooden stake (3 x 3 x 40 cm) inserted in the ground and coated in Oecotack insect trapping glue (Agralan) above ground.

Delta traps were suspended on two bamboo canes (50 cm above the ground) and Unitraps were dug into the ground to the level of the funnel. Four replicates of each trap in each experiment in a randomised block design were spaced >10 m apart and the numbers of male ETB captured recorded weekly.

Investigation of effects of grid on Unitraps for ETB to exclude capture of bees (Apoidea)

The trial was set up in a weed field (*Chenopodium* and *Matricaria*) at NIAB EMR in 2009. Traps were Unitraps with a green or white cross vanes baited with ETB pheromone. The latter were tested with or without a black plastic grid (4 x 5 mm mesh) fitted over the hole of the funnel since white is attractive to bees. Catches of ETB in these traps were compared with those in a green delta trap with a white sticky insert. There were 5 replicates of each treatment in a randomised block design and the traps were spaced >10 m apart. Traps were in place from 11 May - 19 June and from 11 May - 18 August and catches were recorded either weekly or fortnightly.

Effect of cross vane height on catches of ETB and SBW

ETB trials were in the UK on the perimeter of a strawberry crop at NIAB EMR (15 July - 12 August 2013). SBW trials were in a strawberry crop in the north-west area of Norway, Møre and Romsdal County (Lat: 62.697778 north, Long: 7.385278 east) (2 July – 27 August 2013). Traps were baited with the corresponding synthetic pheromone lures.

For the ETB trial, Unitraps had either no cross vane (lid attached directly to bucket), a normal size cross vane (10 cm high), or the cross vane area was doubled by joining up two cross vanes with a 1 ml pipette tip on each corner to secure the vanes (20 cm high). For the SBW test, Unitraps had

either no cross vane, a standard height (10 cm) cross vane or a vane which was half the height (5 cm). All cross vanes were green and the traps stood on the ground and held in place with a wire hoop.

There were 10 replicates of each treatment arranged in a randomised complete block design. Plots were rows with single traps deployed spaced 10 m apart for ETB and 20 m apart for SBW. Counts of SBW and male ETB were made weekly.

Comparison of trap designs for trapping of SBW and ETB

Ten different trap designs were compared to find one effective trap for both SBW and ETB. Experiments were carried out in the UK and Latvia on two occasions, one to coincide with SBW emergence (UK 19 March – 05 July 2012; Latvia 17 May – 19 July 2012) and the second with the ETB main flight period (UK 23-31 July 2012; Latvia 30 July - 31 August 2012).

In the UK, organic strawberry plantations in Hereford (Lat: 52.050051 north, Long:-2.491226 west) were utilised for the SBW experiment and a strawberry crop at NIAB EMR for the ETB test. In Latvia the SBW trial was set up in organic strawberry plantations (Lat: 57.113804 north, Long: 24.530401 east; Lat: 56.807688 north, Long: 24.271681 east; Lat: 56.630664 north, Long: 23.344844 east; Lat: 56.595720 north, Long: 23.272959 east; Lat: 56.921632 north, Long: 23.211209 east) and the ETB trial in a sowing of *Medicago sativa* L. near Vecauce (Lat: 56.595720 north, Long: 23.27295 east).

Trap designs tested are shown in Table 2 and were baited with the synthetic SBW or ETB lures according to the target pest. Traps included Unitraps with different coloured cross vanes, with and without excluders. The effect of attaching the lure inside the bucket rather than to the top of the cross vanes was tested. Traps made of a vertical cylinder of card (25 cm x 10 cm dia) coated with dry glue in different colours were tested as these are supplied commercially for ETB (Agralan), as were the Xlure-RTU sticky trap (12 cm x 3 cm) from Russell IPM. Two simple, homemade traps were

tested: the sticky stake trap and a white plastic strip (45 x 150 x 3 mm) in 45 mm pot containing water + 0.1% Triton X100.

The two experiments (one in the UK and one in Latvia) were set up as a randomised complete block design. There were five replicates of each treatment in each experiment. Because of the area needed for each replicate, experiments were conducted across several plantations, with each considered a replicate with all 10 treatments. Traps were deployed 15-20 m apart along the edges of the fields. The traps were stood on the ground, with foliage around them removed. Sticky traps were set with their bottom edge touching the ground. Unitraps were held in place with a wire hoop and contained water plus a drop of detergent.

Total counts of SBW and male ETB in each trap were made every 2 weeks. Bycatch of other notable insects were recorded into broad taxa and included honeybees, bumblebees, solitary bees and Diptera (>2 mm).

Effect of trap height and habitat on catches of ETB and SBW

The ETB trial was set up in a strawberry plantation (cv. 'Finesse') at NIAB EMR (14 August - 26 September 2011). Treatments were green cross vane Unitraps with water and detergent as a trapping agent. Traps were wired in position at ground level (0 m) or onto the bracing bar of the tunnel (1.25 m from the ground) or onto the centre top ridge pole of tunnel (4 m from the ground). Traps were baited with the synthetic ETB sex pheromone lure. The trial had a randomised complete block design with 5 replicates and the traps were arranged around the edge of the plantation >10 m apart. Trap catches were recorded weekly.

The SBW trials were carried out in raspberry in 2012 at three locations in Denmark, Norway and Switzerland (Table 3). Three habitat types were tested; the crop (raspberry), the boundary of the crop and 50 m into the adjacent forest. Traps were green Unitraps with a white cross vane with a synthetic SBW lure, a pollinator exclusion grid across the funnel opening, and water with detergent as the trapping agent in the bucket. Traps were mounted on poles at three heights above ground

level (0 m, 0.5 m and 1.5 m). There were 9 replicates in Denmark and Norway and 12 in Switzerland. Traps in each habitat were >3 m apart and catches recorded every 7-14 days.

Data analysis

All analyses were carried out using Genstat v 14 (VSN International, 2011). Mean total catches during the experiments were square root transformed, to stabilise the variances, and subjected to analysis of variance (ANOVA). Where significant differences between means were indicated, these were tested for significance by a Least Significant Difference (LSD) test using $P < 0.05$ to indicate significant differences.

For the UK experiments fixed effects were the treatments (e.g. trap design, cross vane height, trap height, habitat etc.) and random effects were the blocks in the randomised experiment designs.

The Latvia and UK data for the trap type experiment to capture SBW or ETB were combined and analysed using ANOVA. The fixed effects were trap type and country and random effects were blocks nested within country and treatments nested within blocks. Data from this experiment was also analysed using a Generalised Linear Mixed Model (GLMM) (Breslow, N.E. & Clayton, D.G., 1993) with the Poisson distribution and a log-link. This was analysed with the same fixed effects and the same random effect model as the ANOVA, above. Where the ANOVA gave significant trap type x country interactions for all variables, this was not the case in the GLMM analyses for any of the variables. This lack of interaction in the GLMM with its log-link implies that the trap type means differed between the two countries by a fixed multiplier which corresponded to the relative abundances in the two countries. To preserve conformity with other analyses the overall means of trap type were reported from the ANOVA analyses.

When analysing the effect of trap height on catches of SBW catches, mean counts over the trapping season for countries (Norway, Denmark, Switzerland) were analysed using ANOVA with trap

height, habitat and country as fixed effects and replicates within country, and height within blocks as random effects. A GLMM analysis was also used but in this case gave similar results to the ANOVA.

Results

Comparison of delta traps and Unitraps for trapping ETB

In Experiment A, catches of ETB were low and there were no significant differences between the trap types tested ($F_{6,18} = 1.22$, $P=0.343$; data not shown). The addition of lambda-cyhalothrin or cypermethrin to the traps did not significantly improve ETB catches.

In Experiment B, there were significantly more male ETB captured in the green cross vane Unitrap compared to the green or clear delta traps ($F_{6,18}=2.16$, $P=0.096$, $l_{sd}=2.768$, Figure 1). Unitraps with white or yellow cross vanes had intermediate trap catches.

Investigation of effects of a grid on Unitraps for ETB to exclude capture of bees (Apoidea)

Using a plastic grid over the hole of the funnel in the white cross vane Unitraps significantly reduced the numbers of captured bumblebees ($F_{4,16} = 6.25$, $P=0.003$, $l_{sd}=0.837$, Figure 2) and honeybees ($F_{4,16}=60.44$, $P<0.001$, $l_{sd}=0.756$, Figure 2). Although the white cross vane Unitraps had fewer ETB if a grid was placed over the funnel hole, this was not significant. The main differences were that the green cross vane Unitrap without a grid captured significantly more male ETB than the green Unitrap with a grid or a green delta trap with a sticky glue insert ($F_{4,16}=7.24$, $P=0.002$, $l_{sd}=1.248$, Figure 2). These results suggested that a grid is not needed if a green cross vane is used in the Unitraps. Green cross vanes reduced the bycatch whilst maintaining a significant catch of male ETB.

Effect of cross vane height on catches of ETB and SBW

In the UK trial, the height of the green cross vane had no effect on the numbers of male ETB trapped in the Unitraps. There was also no effect on bumblebee ($F_{2,18}=1.19$, $P=0.328$, $l_{sd}=0.380$), carabid ($F_{2,18}=0.08$, $P=0.924$, $l_{sd}=0.876$), spider ($F_{2,18}=0.78$, $P=0.474$, $l_{sd}=0.716$), earwig ($F_{2,18}=1.47$, $P=0.257$,

l_{sd}=0.689), large Diptera ($F_{2,18}=0.92$, $P=0.417$, l_{sd}=0.845) or ant numbers ($F_{2,18}=0.10$, $P=0.908$, l_{sd}=0.695). However, in general, captures of some notable groups of invertebrates did increase as the vane height was increased, including the mirid *Calocoris norvegicus* (Gmelin) ($F_{2,18}=11.12$, $P<0.001$, l_{sd}=0.729), slugs ($F_{2,18}=3.58$, $P=0.049$, l_{sd}=1.010), Coccinellidae ($F_{2,18}=5.68$, $P=0.012$, l_{sd}=0.801) and Opilione ($F_{2,18}=9.04$, $P=0.002$, l_{sd}=0.718, Figure 3).

In the trial in Norway, SBW catches were significantly lower in the half height (5 cm) cross vane Unitraps compared to the full height cross vane traps ($F_{2,18}=5.21$, $P=0.016$, l_{sd}=0.681, Figure 4). When the cross vane was removed altogether, intermediate numbers of SBW were captured.

Comparison of trap designs for trapping of SBW and ETB

For the SBW and ETB experiments there was no evidence of a trap type x country interaction in the GLMM analysis for SBW counts, ETB counts, or any of the by-catches (not reported), but there were significant differences between the trap types. Numbers of SBW captured in the Unitraps with white or green cross vanes, even with the 4 x 5 mm excluder grid, ($F_{9,72}=13.42$, $P<0.001$, l_{sd}=0.889, Figure 5A) were significantly higher than in other trap types. The lure placed inside the bucket of the Unitrap, instead of under the lid, captured fewer SBW and was comparable in captures to the sticky stake, the pot trap and the yellow sticky card traps (Figure 5A).

By-catch was also affected by trap design. Higher numbers of bumblebees, in particular, were captured in the Unitraps with the white cross vanes and no, 4 x 5 mm, excluder grid compared to most other traps ($F_{9,72}=10.74$, $P<0.001$, l_{sd}=0.418, Figure 5B). This was also true for honeybees ($F_{9,72}=2.83$, $P=0.007$, l_{sd}=0.583, Figure 5C) and solitary bees ($F_{9,72}=5.15$, $P<0.001$, l_{sd}=0.632, Figure 5D). However significant numbers of solitary bees were also captured in the pot trap and blue and yellow dry sticky glue traps.

Large Diptera (>2 mm) were more abundant on traps which had glue as the trapping agent in comparison to liquid based traps and the Xlure R.T.U floor trap ($F_{9,72}=52.12$, $P<0.001$, l_{sd}=1.986, Figure 5E).

Later in the season, in the ETB trial, more ETB were captured in green or white cross vane Unitraps without the excluder grid compared to other trap types, although small numbers were captured in the pot trap and in the Unitrap with the excluder grid ($F_{9,72}=33.76$, $P<0.001$, $l_{sd}=1.414$, Figure 6A). Placing the pheromone lure inside the bucket of the Unitrap did not increase catches of ETB (Figure 6A).

By-catch, later in the season, included honeybees and large Diptera. Honeybees were more likely to be captured in the Unitraps with the white cross vane and without the 4 x 5 mm excluder grid; small but significant numbers of honeybees were also captured on the blue sticky card ($F_{9,72}=3.57$, $P=0.001$, $l_{sd}=0.598$, Figure 6B). As with the previous study, earlier in the season, by-catches of large Diptera (>2 mm) in the ETB experiment were higher on traps which comprised sticky glue, including the yellow wet and dry cards, the blue card and the sticky stake trap, in comparison to traps which used a liquid as the method of capture ($F_{9,72}=50.09$, $P<0.001$, $l_{sd}=1.333$, Figure 6C).

Effect of trap height and habitat on catches of ETB and SBW

In the UK experiment there were significantly more male ETB caught in traps placed on the ground within the strawberry row (mean 22.2) than in the traps wired to the tunnel bracing bar 1.25 m above ground (mean 1.0) or the ridge pole 4 m above ground (mean 0.2) ($F_{2,8}=29.13$, $P<0.001$, $l_{sd}=1.558$, data not shown).

There were significant interactions between country and habitat ($F_{4,14} = 10.61$, $P<0.001$, $l_{sd} = 0.55$), and habitat and trap height ($F_{4,42}=3.13$, $P=0.024$, $l_{sd}=0.257$). There were no other significant interactions. The numbers of SBW captured in Denmark and Switzerland were lower than those in Norway, particularly in raspberry crops and the crop boundary compared to the forest (Figure 7A). In the boundary and the crop, SBW was more likely to be trapped at 0 m compared to 0.5 or 1.5 m (Figure 7B).

Discussion

In this study we developed and tested traps for SBW and ETB monitoring. The best trap, effectively capturing both species with minimal bycatch, was the green cross vane Unitrap with no excluder grid over the hole to the funnel (Figure 8). This trap gave best capture of the two pest species if deployed on the ground. The white cross vanes previously used to trap SBW (Cross *et al.*, 2006a, b; Wibe *et al.*, 2014) did not improve capture compared to green cross vanes and more bees were attracted. Bees could be excluded by use of a plastic mesh grid over the Unitrap funnel entrance, but this impeded the capture of ETB. Water and detergent were better for trapping ETB than glue, and ETB and SBW have been observed to free themselves from glue traps (pers. obs.). Overall the sticky traps were unsuitable for trapping SBW and ETB because they became contaminated and potentially saturated with other arthropods.

The height of the Unitrap cross vanes did not affect the capture of ETB, but the 10 cm cross vanes were most effective for capture of SBW. However, an increase in cross vane height did increase the capture of some beneficial arthropods including Coccinellidae and Opilione. Trap contaminants, in particular slugs, seemed to increase with vane height and hence the management of slugs through irrigation control and crop management is needed. In addition, the height which the trap is placed affected trapping efficacy. The ETB traps placed near to the ground at crop height captured more males than traps placed higher in the tunnel structure. Likewise, SBW was more likely, in Norway, to be captured on the ground in the forest. This may be because the wild fruits are closer to the ground compared to commercially grown fruits which are tied vertically into post and wire systems with the fruit higher than would be natural.

Although this trapping system is cost effective for monitoring, further improvements and reduction in cost need to be made for mass trapping. An attempt was made to find a low cost, small, trap that could be deployed in large numbers for mass trapping. However, none of the traps, including prototypes, were as effective as the Unitrap with green cross vanes (Figure 8).

An obvious flaw in the trap system tested here is the lack of an attractant for female ETB. However, phenylacetaldehyde and/or (*E*)-cinnamaldehyde (Koczor *et al.*, 2012) could be added to

the sex pheromone to increase catches of female ETB. An alternative method to the drowning solution, which needs to be emptied and topped up on a monthly basis, would make the trap easier to maintain. There is the potential to incorporate an insecticide (Navarro-Llopis *et al.*, 2014) or biological control agent onto the inside of the trap. This would be a lure and kill system where the insect would enter and then die either inside the trap or after leaving.

In addition, the lures also need maintenance. SBW lures last at least two months under field conditions (Cross *et al.*, 2006a), but the longevity of the ETB lures is approximately four weeks (Fountain *et al.*, 2014). It would be beneficial to increase the longevity of these lures and/or improve the deployment and replacement of lures in the traps. Finally, the traps were often placed in the leg row of the strawberry crop to avoid disturbance by spray machinery and fruit pickers, but this made them difficult to access on a regular basis (see Figure 8).

In this study we did not test lures for both species in the trap at the same time. Further studies are required to ensure that the components of the two lures remain attractive to the pest species when placed together. It is considered unlikely there would be any interaction in view of the very different chemical structures of the pheromone components. A combined trap would save time in monitoring and push the cost-benefit ratio of mass trapping in the right direction. There is a real potential to mass trap (Fountain *et al.*, 2015) or lure and kill ETB and SBW in strawberry crops. Future research should concentrate on trap optimization, ease of use and economics of deploying and maintaining these systems for reducing fruit damage in strawberry crops.

Acknowledgements

This work was supported by the “Softpest Multitrap” project, funded by FP7-ERA-Net CORE Organic II and Hort LINK project HL0184/PC, SF 276 “Pheromone technology for management of capsid pests to reduce pesticide use in horticultural crops” funded by the UK AHDB and Defra. We would like to thank the numerous fruit growers who allowed access to their crops for the trap testing, the numerous technical assistants and students who helped in trap deployment and insect counts.

References

- Aasen, S.S. & Trandem, N. (2006). Strawberry blossom weevil *Anthonomus rubi* Herbst (Col.: Curculionidae): relationships between bud damage, weevil density, insecticide use, and yield. – *Journal of Pesticide Science*, **79**, 169-174.
- Abbes, K., Harbi, A. & Chermiti, B. (2012). Comparative study of 2 protection strategies against *Tuta absoluta* (Meyrick) in late open field tomato crops in Tunisia. *EPPO Bulletin*, **42**, 297-304.
- Bichão, H., Borg-Karlson, A.-K., Araujo, J. & Mustaparta, H. (2005a). Five types of olfactory receptor neurons in the strawberry blossom weevil *Anthonomus rubi*: selective responses to inducible host-plant volatiles. *Chemical Senses*, **30**, 153-170.
- Bichão, H., Borg-Karlson, A.-K., Wibe, A., Araujo, J. & Mustaparta, H. (2005b). Molecular receptive ranges of olfactory receptor neurones responding selectively to terpenoids, aliphatic green leaf volatiles and aromatic compounds, in the strawberry blossom weevil *Anthonomus rubi*. *Chemoecology*, **15**, 211-226.
- Blackmer, J. L., Byers, J. A. & Rodriguez-Saona, C. (2008). Evaluation of color traps for monitoring *Lygus* spp.: Design, placement, height, time of day, and non-target effects. *Crop Protection*, **27**, 171-181.
- Breslow, N.E. & Clayton, D.G. (1993). Approximate inference in generalized linear mixed models. *Journal of the American Statistical Association*, **88**, 421, 9-25.
- Cressey, D. (2015). Bee studies stir up pesticide debate. *Nature*, **520**, 416.
- Croft, B.A. & Brown, A.W.A. (1975). Responses of arthropod natural enemies to insecticides. *Annual Review of Entomology*, **20**, 285-335.
- Cross, J., Fountain, M., & Hall, D. (2011). Management of European tarnished plant bug in late season strawberries. *IOBC/WPRS Bulletin*, **70**, 170.

- 400 Cross, J.V., Hesketh, H., Jay, C.N., Hall, D.R., Innocenzi, P.J., Farman, D.I. & Burgess C.M. (2006a).
 401 Exploiting the aggregation pheromone of strawberry blossom weevil *Anthonomus rubi* Herbst
 402 (Coleoptera: Curculionidae): Part 1. Development of lure and trap. *Crop Protection* **25**, 144-154.
- 403 Cross, J.V., Hall, D.R., Innocenzi, P.J., Hesketh, H., Jay, C.N. & Burgess, C.M. (2006b). Exploiting the
 404 aggregation pheromone of strawberry blossom weevil *Anthonomus rubi* (Col.: Curculionidae):
 405 Part 2. Pest monitoring and control. *Crop Protection*, **25**, 155-166.
- 406 Dembilio, O. & Jaques, J.A. (2015). Biology and management of red palm weevil. In Sustainable Pest
 407 Management in Date Palm: Current Status and Emerging Challenges (Eds W. Wakil, J.R. Faleiro &
 408 T.A. Miller). Springer, Switzerland, pp 13-36.
- 409 El-Sayed, A.M, Suckling, D.M, Wearing, C.H. & Byers, J.A. (2006). Potential of mass trapping for long-
 410 term pest management and eradication of invasive species. *Journal of Economic Entomology*, **99**,
 411 1550-64.
- 412 European Food Safety Authority (2015). The 2013 European Union report on pesticide residues in
 413 food. EFSA Journal;13(3):4038, 169 pp. doi:10.2903/j.efsa.2015.4038
- 414 Faccoli, M. & Stergulc, F. (2008). Damage reduction and performance of mass trapping devices for
 415 forest protection against the spruce bark beetle, *Ips typographus* (Coleoptera Curculionidae
 416 Scolytinae). *Annals of Forest Science*, **65**, 309p1-309p9.
- 417 Fitzgerald J. & Jay C. (2011). Chemical control of the European tarnished plant bug, *Lygus*
 418 *rugulipennis*, on strawberry in the UK. *Crop Protection*, **30**, 1178-1183.
- 419 Fountain, M., Cross, J., Jaastad, G., Farman, D. & Hall, D. (2010). Developing an effective trap and
 420 lure to monitor *Lygus rugulipennis*. *IOBC/wprs Bulletin*, **54**, 47–51.
- 421 Fountain, M., Cross, J., Jaastad, G. & Hall, D. (2008). Some preliminary investigations into the sex
 422 pheromones of mirid soft fruit pests. *IOBC/WPRS Bulletin*, **39**, 33-40.
- 423 Fountain, M.T., Jåstad, G., Hall, D., Douglas, P., Farman, D. & Cross J.V. (2014). Further studies on sex
 424 pheromones of female *Lygus* and related bugs: Development of effective lures and investigation
 425 of species-specificity. *Journal of Chemical Ecology*, **40**, 71-83.

- 426 Fountain, M.T., Cross, J.V., Farman, D. & Hall, D. (2011) Using pheromones to monitor *Lygus*
 427 populations in fruit crops in the UK. *Pheromones and other semio-chemicals IOBC/wprs Bulletin*,
 428 **72**, 7-13.
- 429 Fountain, M.T., Shaw, B., Trandem, N., Storberget, S., Baroffio, C., Ralle, B., Rendina, P., Richoz, P.,
 430 Sigsgaard, L., Borg-Karlson, A.-K., Hall, D., Cross, J.V. & Wibe, A. (2015). The potential for mass
 431 trapping *Lygus rugulipennis* and *Anthonomus rubi*; trap design and efficacy. *Integrated protection*
 432 *of fruit crops Subgroup "Soft Fruits" IOBC-WPRS Bulletin*, **109**, 95-97.
- 433 Frati, F., Chamberlain, K., Birkett, M., Dufour, S., Mayon, P., Woodcock, C., Wadhams, L., Pickett, J.,
 434 Salerno, G., Conti, E. & Bin, F. 2009. *Vicia faba* – *Lygus rugulipennis* interactions: induced plant
 435 volatiles and sex pheromone enhancement. *Journal of Chemical Ecology*, **35**, 201-208.
- 436 Haghani, M., Negahban, M., Sedaratian-Jahromi, A. & Ghane-Jahromi, M. (2016). Monitoring of an
 437 Iranian population of *Grapholita funebrana* Treitschke, 1835 (Lepidoptera: Tortricidae) using sex
 438 pheromone traps: An applicable procedure for sustainable management. *Entomofauna*, **37**, 241-
 439 252.
- 440 Hillocks, R.J. (2012). Farming with fewer pesticides: EU Pesticide Review and resulting challenges for
 441 UK agriculture. *Crop Protection*, **31**, 85-93.
- 442 Hillocks, R.J. (2013). Impact of EU Pesticide Reduction Strategy and implications for crop protection
 443 in the UK and the rest of Europe. *Outlooks on Pest Management*, **24**, 206-209.
- 444 Holopainen, J.K., Raiskio, S., Wulff, A. & Tiilikkala K. (2001). Blue sticky traps are more efficient for
 445 the monitoring of *Lygus rugulipennis* (Heteroptera, Miridae) than yellow sticky traps. *Agricultural*
 446 *and food science in Finland*, **10**, 277-284.
- 447 Innocenzi, P.J., Hall, D., Cross, J.V. & Hesketh, H. (2005). Attraction of male European tarnished plant
 448 bug, *Lygus rugulipennis* to components of the female sex pheromone in the field. *Journal of*
 449 *Chemical Ecology*, **31**, 1401-1413.

- 450 Innocenzi, P.J., Hall, D.R. & Cross, J.V., (2001). Components of male aggregation pheromone of
 451 strawberry blossom weevil, *Anthonomus rubi* Herbst. (Coleoptera : Curculionidae). *Journal of*
 452 *Chemical Ecology*, **27**, 1203-1218.
- 453 Innocenzi, P.J., Hall, D.R., Cross, J.V., Masuh, H., Phythian. S.J., Chittamaru, S. & Guarino, S. (2004).
 454 Investigation of long-range female sex pheromone of the European tarnished plant bug, *Lygus*
 455 *rugulipennis*: Chemical, electrophysiological, and field studies. *Journal of Chemical Ecology*, **30**,
 456 1509-1529.
- 457 Jay, C., Cross, J. & Burgess, C. (2008). Severe damage by *Anthonomus rubi* populations in the UK.
 458 *IOBC/WPRS Bulletin*, **39**, 131-136.
- 459 Koczor, S., Vuts, J. & Tóth, M. (2012). Attraction of *Lygus rugulipennis* and *Adelphocoris lineolatus* to
 460 synthetic floral odour compounds in field experiments in Hungary. *Journal of Pest Science*, **85**,
 461 239-245.
- 462 Mafra-Neto, A. & Habib, M. (2003). Evidence that mass trapping suppresses pink bollworm
 463 populations in cotton fields. *Entomologia Experimentalis et Applicata*, **81**, 315–323.
- 464 Mwatawala, M.W., Mziray, H., Malebo, H. & De Meyer, M. (2015). Guiding farmers' choice for an
 465 integrated pest management program against the invasive *Bactrocera dorsalis* Hendel (Diptera:
 466 Tephritidae) in mango orchards in Tanzania. *Crop Protection*, **76**, 103-107.
- 467 Navarro-Llopis, V., Primo, J. & Vacas, S. (2015). Bait station devices can improve mass trapping
 468 performance for the control of the Mediterranean fruit fly. *Pest Management Science*, **71**, 923–
 469 927.
- 470 Oehlschlager, A. C., Chinchilla, C., Castillo, G., & Gonzalez, L. (2002). Control of red ring disease by
 471 mass trapping of *Rhynchophorus palmarum* (Coleoptera: Curculionidae). *Florida. Entomologist*,
 472 **85**, 507-513.
- 473 Renkema, J.M., Buitenhuis, R. & Hallett, R.H. (2014). Optimizing trap design and trapping protocols
 474 for *Drosophila suzukii* (Diptera: Drosophilidae). *Journal of Economic Entomology*, **107**, 2107-2118.

- 475 Singh, N., Wang, C. & Cooper, R. (2013). Effect of trap design, chemical lure, carbon dioxide release
 476 rate, and source of carbon dioxide on efficacy of bed bug monitors. *Journal of Economic*
 477 *Entomology*, **106**, 1802-1811.
- 478 Switzer, P.V., Enstrom, P.C. & Schoenick, C.A. (2009). Behavioural explanations underlying the lack of
 479 trap effectiveness for small-scale management of Japanese beetles. *Journal of Economic*
 480 *Entomology*, **102**, 934-940.
- 481 Teixeira, L.A.F., Mason, K. & Isaacs R. (2009). Control of grape berry moth (Lepidoptera: Tortricidae)
 482 in relation to oviposition phenology. *Journal of Economic Entomology*, **102**, 692-698.
- 483 VSN Internation (2011). GenStat for Windows 14th Edition. VSN International, Hemel Hempstead,
 484 UK. Web page: GenStat.co.uk
- 485 Walton, V.M., Daane, K.M. & Pringle, K.L. (2004). Monitoring *Planococcus ficus* in South African
 486 vineyards with sex pheromone-baited traps. *Crop Protection*, **23**, 1089-1096.
- 487 Wibe, A., Borg-Karlson, A.-K., Cross, J. & Sigsgaard, L. (2011). Effective trapping of the strawberry
 488 blossom weevil, *Anthonomus rubi*. *IOBC/WPRS Bulletin*, **70**, 24.
- 489 Wibe, A., Borg-Karlson, A.-K., Cross, J., Bichao, H., Fountain M., Liblikase, I. & Sigsgaard, L. (2014).
 490 Combining 1,4-dimethoxybenzene, the major flower volatile of wild strawberry *Fragaria vesca*,
 491 with the aggregation pheromone of the strawberry blossom weevil *Anthonomus rubi* improves
 492 attraction. *Crop Protection*, **64**, 122-128.
- 493 Witzgall, P., Kirsch, P. & Cork, A. (2010). Sex pheromones and their impact on pest management.
 494 *Journal of Chemical Ecology*, **36**, 80-100.

Table 1 Comparison of traps for capturing male ETB using virgin female ETB as bait in UK.

Trap	Capture device
<i>Experiment A (27 June – 11 July 2008)</i>	
Green delta	Wet glue sticky insert
Green delta	Dry glue sticky insert
Green delta	Wet glue sticky insert + cypermethrin
Green delta	Wet glue sticky insert + Oecotack
Unitrap (white cross vane)	Correx card + water with detergent
Unitrap (lambda-cyhalothrin white cross vane)	Water + detergent
Clear delta	Wet glue sticky insert
<i>Experiment B (27 August – 1 September 2008)</i>	
Green delta	Wet glue sticky insert + Oecotack
Clear delta	Wet glue sticky insert + Oecotack
Agrisense vane funneled (white cross vanes)	Water + detergent
Unitrap (green cross vanes)	Water + detergent
Unitrap (white cross vanes)	Water + detergent
Unitrap (yellow cross vanes)	Water + detergent
Sticky stake trap	Oecotack

495

496

Table 2 Trap designs tested for trapping SBW and ETB in the UK and Latvia. Traps were baited with synthetic SBW or ETB lures.

Trap design	Killing agent	Position of lure	Source of trap
Unitrap: White cross vane with excluder-grid (4x5 mm)	Water + 0.1% Triton X100	Top of cross vane	Agralan
Unitrap: White cross vane with excluder-grid (4x5 mm)	Water + 0.1% Triton X100	Inside bucket attached to funnel	Agralan
Unitrap: White cross vane without excluder-grid	Water + 0.1% Triton X100	Top of cross vane	Agralan
Unitrap: Green cross vane without excluder-grid	Water + 0.1% Triton X100	Top of cross vane	Agralan
Sticky stake trap (2.5 x 2.5 x 50 cm)	Oecotack wet glue	Top of stake	NIAB EMR
Yellow card cylinder (25 x 10 cm)	Coated in wet glue	Top of cylinder	Agralan
Blue card cylinder (25 x 10 cm)	Coated in dry glue	Top of cylinder	Agralan
Yellow card cylinder (25 x 10 cm)	Coated in dry glue	Top of cylinder	Agralan
White plastic strip (45 x 150 x 3 mm) in 45 mm pot	Water + 0.1% Triton X100	Top of strip	NIAB EMR
Xlure R.T.U floor trap	Oecotack sticky insert inside	Inside, in middle on base	Russell

Table 3 Location and age of raspberry plantations used for testing the effect of trap height on catches of SBW (CH Switzerland, NO Norway, DK Denmark).

Country	Location	Coordinates		Production type	Plantation year
		Latitude	Longitude east		
CH	Bruson, Valais	46.0037	7.3191	Open	2005
CH	Nendaz, Valais	46.1834	7.2942	Open	2005
CH	St Sébastien, Valais	46.198	7.31306	Open	2004
NO	Skåla, Molde	62.6953	07.3769	Protected	2010
NO	Skjønby	60.8278	10.7970	Open	2008
NO	Torp	59.6677	10.6912	Open	2010
DK	Gyrstinge	55.4770	11.6830	Open	2010
DK	Eggeslevmagle	55.2843	11.3300	Protected	2010
DK	Kildebrønde	55.6033	12.2650	Protected	2010

Figure 8 Green cross vane Unitrap anchored into place on the ground in a strawberry crop. The pheromone dispensers are deployed in the cage inserted into the lid.

Figure(s)1




Figure 1 Mean numbers (\pm S.E.) of male ETB captured in different traps (Table 1) baited with live female ETB in the UK (27 Aug – 1 Sept 2008; means with the same letter are not significantly different, $P > 0.05$).




Figure 2 Mean catches (\pm S.E.) of bumblebees, honeybees and male ETB in green Unitraps with white or green coloured cross vanes, with and without a grid at the entrance to the funnel in comparison to a green Delta trap (UK, 11 May - 19 June and 11 May - 18 August; means with different letters for each species are significantly different , $P < 0.05$).




Figure 3 Mean numbers (\pm S.E.) of invertebrate bycatch captured in green Unitraps with either no cross vane, a single height (10 cm) or double height (20 cm) green cross vane (UK, 15 July - 12 August 2013; means with different letters for each invertebrate group are significantly different, $P < 0.05$).




Figure 4 Mean numbers (\pm S.E.) of SBW captured in green Unitraps with either no cross vane, a full height (10 cm) or half height (5 cm) white cross vane (Norway, 16 July – 27 August 2013; means with different letters are significantly different , $P < 0.05$).




Figure 5 Mean numbers (\pm S.E.) of A) SBW, B) bumblebees, C) honeybees D) solitary bees and E) large Diptera in ten trap designs (Table 2). Data from the locations pooled by LSD test after ANOVA on square root transformed data. Means with different letters are significantly different. The arithmetic means are presented here, whereas the means of the square-root transformed counts are grouped, hence there may be apparent discrepancies in the groupings.




Figure 6 Mean numbers (\pm S.E.) of A) ETB, B) honeybees and C) large Diptera in ten trap designs (Table 2). Data from the locations pooled by LSD test after ANOVA on square root transformed data. Means with different letters are significantly different. There were no significant differences in the numbers of honeybees in traps in the UK. The arithmetic means are presented here, whereas the means of the square-root transformed counts are grouped hence there may be some discrepancies in the groupings.




Figure 7 Mean number (\pm S.E.) of SBW per trap in A) the boundary, crop (raspberry) or forest in the 3 countries (Denmark, Norway and Switzerland) and B) at different heights (0, 0.5 and 1.5 m) in the 3 habitat types (boundary, crop and forest). Data from 9 (Denmark and Norway) or 12 locations (Switzerland) locations in 3 countries pooled by LSD test after ANOVA on square root transformed data. Means with different letters are significantly different.

Figure(s)8
[Click here to download high resolution image](#)

